

This Kentucky collection provides another location outside of the Coastal Plain, the primary area of distribution, and is the first report from the Interior Low Plateaus. While perhaps a recent introduction, the population size suggests an extensive seed source from previous growth in the same area even though the depression is dry enough for cultivation in some years.

Voucher specimens of *Chester* 83-287 (Kentucky. Christian Co.: 18 Aug 1983) have been deposited in APSU, KNK, SMU, and VDB.—*Edward W. Chester and Kevin Souza, Dept. of Biology, Austin Peay State University, Clarksville, TN 37040, U.S.A.*

REFERENCES

- BRANSON, B. A., D. F. HARKER, J. M. BASKIN, M. E. MEDLEY, D. L. BATCH, M. L. WARREN, W. H. DAVIS, W. C. HOUTCOOPER, B. MONROE, L. R. PHILLIPPE, and P. CUPP. 1981. Endangered, threatened, and rare animals and plants of Kentucky. *Trans. Kentucky. Acad. Sci.* 42:77–89.
- CURREY, M. G., and C. M. ALLEN. 1973. Alismataceae of Louisiana: taxonomy, distribution, and field key. *Proc. Louisiana Acad. Sci.* 36:88–95.
- FASSETT, N. C. 1955. *Echinodorus* in the American tropics. *Rhodora* 57:133–156, 174–188, 202–212.
- HAYNES, R. R. 1980. Aquatic and marsh plants of Alabama. I. Alismatidae. *Castanea* 45:31–50.
- LEONARD, S. W. 1972. New records and notes on the flora of the Carolinas. *J. Elisha Mitchell Sci. Soc.* 88: 265–266.
- LIPSCOMB, B. 1977. *Echinodorus tenellus* var. *parvulus* (Alismataceae) in the Ozarks of Arkansas. *Castanea* 42:254–255.
- MOHLENBROCK, R. H. 1970. The illustrated flora of Illinois, flowering rush to rushes. Southern Illinois University Press, Carbondale. 272 p.
- ROGERS, G. K. 1983. The genera of Alismataceae in the southeastern United States. *J. Arnold Arbor.* 64:383–420.

CAREX ROSEA (CYPERACEAE), *TRIFOLIUM LAPPACEUM* (FABACEAE) AND *AIRA CARYOPHYLLEA* (POACEAE) NEW TO TEXAS—*CAREX ROSEA* Schkuhr is a woodland sedge ranging from Canada south to Louisiana and Oklahoma (Fernald, 1950; Waterfall, 1966). A collection of this species (*Brown* 4805 SMU) in April of 1980 is apparently the first reported for the state. The specimen was collected on a moist wooded seepage slope within Eisenhower Park in northeast Harris Co. near San Jacinto River south of the dam at Lake Houston.

TRIFOLIUM LAPPACEUM L., an introduced clover, was collected in May of 1979 in Harris Co. from a yard adjacent to Kirby Drive in Taylor Lake Village (*Brown* 4370; SMU, TEX) and in Hardin Co. from the roadside of U. S. Hwy. 69 ca 4 miles north of the intersection with Hwy. 420 (*Brown* 4475 SMU).

AIRA CARYOPHYLLEA L. was found in 1981 on sandy soil within the Little Thicket Nature Sanctuary ca 2 miles south of Evergreen on Hwy. 945 in San Jacinto Co. (*Brown and Peterson* 5307 SMU, TAES). This site is ca

170 miles west of the nearest recorded locality in Evangeline Parish, Louisiana (Allen, 1981).—*Larry E. Brown, Department of Biology, Houston Community College, Houston, TX 77007, U.S.A.; Charles D. Peterson, R.A. Vines Science Center, Houston, TX 77055, U.S.A.*

REFERENCES

- ALLEN, C. M. 1981. Grasses of Louisiana. The University of Southwestern Louisiana, Lafayette.
- CORRELL, D. S. & M. C. JOHNSTON. 1970. Manual of the vascular plants of Texas. Texas Research Foundation, Renner.
- FERNALD, M. L. 1940. Gray's manual, 8th ed. Amer. Book Co., N. Y.
- HERMAN, F. J. 1953. A botanical synopsis of the cultivated clovers (*Trifolium*). U. S. Dep. Agric. Monogr. 22.
- WATERFALL, U. T. 1966. Keys to the flora of Oklahoma, 3rd ed. Student Union Book Store. Oklahoma State University, Stillwater.